



VERIFICATION OF A TRANSLATION

I, the below named translator, hereby declare that:

My name and post office address are as stated below:

That I am knowledgeable in the English language and in the language in which the below identified application was filed, and that I believe the English translation of the Japanese Patent Application No. 220060/1998 is a true and complete translation of the above-identified Japanese Patent Application as filed.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Date

6th day of September, 2002

Full name of the translator Kiyoshi MURAKAMI

Signature of the translator

Kiyoshi Murakami

Post Office Address c/o YUASA AND HARA, Section 206,
New Ohtemachi Bldg., 2-1, Ohtemachi
2-chome, Chiyoda-ku, Tokyo, JAPAN

RECEIVED

NOV 19 2002

TECH CENTER 1600/2000

PATENT OFFICE
JAPANESE GOVERNMENT

This is to certify that the annexed is a true copy of the following application as filed with this Office.

Date of Application: August 4, 1998

Application Number : Patent Application
No. 220060/1998

Applicant(s): JAPAN TOBACCO INC.

Dated this 19th day of August, 1999

Commissioner, Takeshi ISAYAMA (Seal)
Patent Office

ISSUANCE No. 11-3058016

[Name of Document] Patent Application

[Case No] 980687

[Filing Date] August 4, 1998

[To] Commissioner, Patent Office

[International Patent Classification]

C12N

[Title of the Invention] MUTANT BARNASE GENE AND TRANSGENIC
PLANT TRANSFORMED BY SAID GENE

[Number of the Claims] 9

[Inventor]

[Address] c/o Japan Tobacco Inc., Plant Breeding
and Genetics Research Laboratory of 700,
Higashibara, Toyoda-cho, Iwata-gun,
Shizuoka

[Name] Kazuyuki HAMADA

[Inventor]

[Address] c/o Japan Tobacco Inc., Plant Breeding
and Genetics Research Laboratory of 700,
Higashibara, Toyoda-cho, Iwata-gun,
Shizuoka

[Name] Fumio NAKAKIDO

[Applicant for Patent]

[Identification No.] 000004569

[Appellation] JAPAN TOBACCO INC.

[Proxy]

[Identification No.] 100089705

[Address] YUASA AND HARA, Section 206,
New Ohtemachi Bldg., 2-1,
Ohtemachi 2-chome, Chiyoda-ku, Tokyo

[Name] Ichio SHAMOTO

[Telephone] 03-3270-6641

[Proxy]

[Identification No.] 100071124

[Patent Attorney]

[Name]	Shosuke IMAI	
[Proxy]		
[Identification No.]	100076691	
[Patent Attorney]		
[Name]	Chuji MASUI	
[Proxy]		
[Identification No.]	100075236	
[Patent Attorney]		
[Name]	Tadahiko KURITA	
[Proxy]		
[Identification No.]	100075270	
[Patent Attorney]		
[Name]	Yasushi KOBAYASHI	
[Proxy]		
[Identification No.]	100092886	
[Patent Attorney]		
[Name]	Kiyoshi MURAKAMI	
[Fees]		
[Page No. of Books]	051806	
[Amount]	¥21,000.-	
[List of the Documents]		
[Item]	Specification	1
[Item]	Abstract	1
[Proof, requested or not]	requested	

[Name of Document] Specification

[Title of the Invention]

MUTANT BARNASE GENE AND TRANSGENIC
PLANT TRANSFORMED BY SAID GENE

[Claims]

[Claim 1] A mutant barnase gene having one or some mutations at least in part of the DNA sequence of barnase gene, wherein said mutant barnase gene is capable, when anther-specifically expressed in a plant, of making said plant substantially male sterile.

[Claim 2] A gene which is derived from a DNA sequence encoding the same amino acid sequence as shown in SEQ ID NO:1 by substitution, deletion, insertion or addition of one to several nucleotides in said DNA sequence, wherein said gene is capable of encoding a protein which makes a plant substantially male sterile when expressed anther-specifically in said plant.

[Claim 3] A gene comprising a DNA sequence which is derived from a DNA sequence encoding the same amino acid sequence as shown in SEQ ID NO:1 by substitution, deletion, insertion or addition of one to several nucleotides in the latter DNA sequence, and a promoter located upstream of said DNA sequence for allowing an anther-specific expression, wherein said gene is capable of making a plant substantially male sterile when introduced into the genome of said plant.

[Claim 4] A gene represented by SEQ ID NO:3 which encodes a protein capable of making a plant substantially

male sterile when expressed anther-specifically in said plant.

[Claim 5] A gene comprising the sequence represented by SEQ ID NO:3 and a promoter located upstream of said sequence for allowing an anther-specific expression, wherein said gene is capable of making a plant substantially male sterile when introduced into the genome of said plant.

[Claim 6] A recombinant vector which contains a gene as claimed in any of Claims 1 to 5 and expresses said gene in a host plant.

[Claim 7] A method of making a plant male sterile which comprises transforming said plant by a mutant barnase gene as claimed in any of Claims 1 to 5 and allowing said mutant barnase gene to be expressed anther-specifically.

[Claim 8] The method as claimed in Claim 7, wherein said plant is transformed by said mutant barnase gene by integrating said gene into the genome of said plant.

[Claim 9] A transgenic plant wherein a gene as claimed in any of Claims 1 to 5 has been introduced.

[Detailed Description of the Invention]

[0001]

[Technical Field of the Invention]

This invention relates to a mutant barnase gene which makes it possible to efficiently yield a male sterile transgenic plant when expressed in a specific site of a plant, in particular, anther-specifically. The

present invention further relates to a recombinant vector capable of expressing the mutant barnase gene of the invention in host cells, a plant transformed by this vector and a method for constructing a transgenic plant.

[0002]

[Prior Art]

Barnase is an RNase originating in *Bacillus amyloliquefaciens* (S. Nishimura and M. Nomura, Biochem. Biophys. Acta 30, 430-431:1958; R.W. Hartley, J. Mol. Biol., 202, 913-915:1988). This enzyme has 110 amino acid residues and hydrolyzes RNA. When expressed in cells, this enzyme inhibits the functions of the cells as a result of its potent RNase activity and thus causes cell death in many cases. By using this characteristic, it is therefore expected that the function of the specific site can be selectively controlled by expressing the barnase gene in a specific site of a plant.

[0003]

PCT International Publication WO89/10396 discloses a technique whereby a male sterile plant is obtained by constructing a male sterility gene by ligating the above-described barnase gene to the downstream of an anther tapetal cell-specific expression promoter and introducing the thus obtained gene into a plant. Male sterilization techniques are of great value in efficiently developing an F1 hybrid variety.

[0004]

When the barnase gene is employed as a male

sterility gene, however, it is frequently observed that resulting male sterile transgenic plants exhibit unfavorable characteristics. PCT International Publication WO96/26283 refers to this problem in rice. It is also reported that similar phenomena are observed not only in rice but in lettuce (Scientia Horticulturae 55, 125-139:1993; Arlette Reymaerts, Hilde Van de Wiele, Greta De Sutter, Jan Janssens: Engineered genes for fertility control and their application in hybrid seed production). According to this report, a plant with depressed activity was constructed by introducing a male sterility gene comprising a tobacco anther-specific promoter (TA29) and a barnase gene into lettuce.

[0005]

Although reasons for these phenomena have not been accurately clarified so far, it is assumed that the so-called "position effect" of the site in which the gene is transferred may account for the mechanism (PCT International Publication WO96/26283). More specifically, a desired male sterile plant should be constructed if the male sterility gene is expressed exclusively in the target site, i.e., anthers. However, there is a possibility that the barnase might be expressed also in tissues other than anthers although in a very small dose under the action of expression regulators, for example, an endogenous enhancer, existing in the vicinity of the gene transfer site. In such a case, the unfavorable characteristics as described above can appear because barnase has a strong enzymatic

activity.

[0006]

To overcome this problem, the method disclosed in PCT International Publication WO96/26283 exploits the character of cauliflower mosaic virus 35S promoter (hereinafter referred to as CaMV35S promoter) of being expressed potently in tissues other than anthers. Namely, barstar, i.e., an inhibitory protein against barnase is employed therein. The barstar gene ligated to a CaMV35S promoter is transferred into a plant simultaneously with a barnase gene and then the barstar gene is constitutively expressed in tissues other than the anthers, thereby eliminating the effects of the barnase in tissues other than the anthers. However, it is necessary in this method to transfer not only a barnase gene but also a barstar gene, and hence, the problem is that the application of this technique to the breeding of F1 varieties of rice or corn seeds may give rise to a gene silencing. "Gene silencing" is a phenomenon wherein the expression of a gene is inhibited when plural copies of the foreign gene are introduced. It is known that this problem frequently occurs when the expression of a foreign gene is effected by a 35S promoter, though the detailed mechanism thereof has not been clarified yet (R.B. Flavell, Proc. Natl. Acad. Sci. USA 91, 3490-3496:1994; J. Finnegan, Bio. Technology 12, 883-888:1994; M.A. Matzke and A.J.M. Matzke, Plant Physiol., 107, 679-685:1995). Since, in rice and corn, a barstar gene is used as a "fertility-restoring gene" in

the pollen parent (father) so as to allow the pollen in the F1 generation to restore the fertility (C. Mariani, et al., Nature 357, 384-387:1992), if an MS plant (mother) is constructed by the method with the use of a barstar gene as reported in WO96/26283, the F1 plant has plural copies of the barstar gene. In such a case, there is a possibility that the expression of the gene may be inhibited due to the gene silencing.

[0007]

[Problems to be Solved by the Invention]

The present invention provides a method for constructing a male sterile plant by using a barnase gene without resort to a barstar gene.

[0008]

The present invention further provides a mutant barnase gene to be used in the above method and a process for producing the same.

[0009]

[Detailed Description of the Invention]

In the present invention, the DNA sequence of barnase gene (R.W. Hartley, J. Mol. Biol. 202, 913-915:1988) is mutated at least in part and then the thus obtained mutant barnase gene is anther-specifically expressed in a plant so as to make the plant substantially male sterile without any substantially disadvantageous effect on the tissues other than anthers.

[0010]

The mutation can be performed by a known method such

as site-specific mutagenesis, deletion of a fragment by using restriction enzymes or the low fidelity PCR method. Among all, it is preferred to use the low fidelity PCR method. This method is described in detail in D. Leung, E. Chen and D. Goedda, Technique 1, 11-15:1989; Y.Z. Xiaoping and R.H. Ebright, Nucleic Acid Res. 19, 6052:1991; G.C. Rice et al., Proc. Natl. Acad. Sci. USA 89, 5467-5471: 1992. By using this technique, PCR is carried out under such conditions as to induce some errors during the amplification reaction. Thus, random mutations can be introduced into the target DNA fragment.

[0011]

Primers to be used in the low fidelity PCR method in the present invention are selected as in the conventional PCR method. It is preferred that each of these primers has a similar number of nucleotides as in the conventional PCR method.

[0012]

The present inventors performed PCR by using a DNA containing the sequence represented by SEQ ID NO:1 as a template with the use of a combination of primer 1 (5'-CGTTCGGCTC GATGGTACCG GTTATCAACA CGTTTGA-3': SEQ ID NO:6) and primer 2 (5'-CCTCTAGATT ATCTGATTTT TGTAAGGTC TGATAATG-3': SEQ ID NO:7) under such conditions as to induce errors. Thus, a mutant barnase gene having the DNA sequence represented by SEQ ID NO:3 was isolated.

[0013]

The sequence represented by SEQ ID NO:1 employed

herein is the sequence of the coding region of the barnase gene contained in the known plasmid pVE108 (WO92/13956). This sequence corresponds to the wild type barnase gene from which the unnecessary moiety corresponding to the secretion signal in the N-terminal side has been deleted.

[0014]

It is also possible to acquire different mutant barnase genes by a similar method.

[0015]

The PCR amplification product is cloned in a host such as *Escherichia coli* and clones containing a mutant barnase gene is isolated. In this cloning, clones having a mutant barnase gene may be screened by assaying the RNase activity expressed by the gene. However, it is advantageous to screen the clone by the method consisting of the following two steps by taking advantage of the fact that the RNase activity of barnase affects the growth of *E. coli*.

[0016]

In the first step, a plasmid is prepared by using the mutant barnase gene obtained above and then *E. coli* is transformed thereby. The growth of the *E. coli* transformant thus obtained will be inhibited by the activity of the mutant barnase. Based on this fact, a colony growing slowly (i.e., a small colony), compared with *E. coli* having a control vector free from any barnase gene, is selected. It is expected that the thus selected *E. coli* strain will contain the mutant barnase gene. To

confirm that the inhibition of the growth of *E. coli* is due to the expression of the mutant barnase gene, the second step is then carried out.

[0017]

In the second step, a barstar gene is employed. As described above, barstar is an inhibitory protein against barnase. In *E. coli* which expresses barstar, the enzymatic activity of barnase is inhibited and thus the degradation of mRNA, which otherwise takes place in the presence of barnase, can be inhibited. Therefore if, *E. coli* which expresses a barstar gene is transformed with a barnase gene, the growth will not be inhibited. Accordingly, it is expected that the thus obtained *E. coli* transformant will produce little difference in growth rate from the transformant constructed by the control plasmid which is free from any barnase gene and, therefore, these transformants will produce little difference in colony size too. Based on this principle, a plasmid is prepared from the *E. coli* strain selected in the first step. The plasmid is then used to transform another *E. coli* strain, in which the barstar gene is constitutively expressed. Thus, mutant barnase gene-containing colonies having almost the same size as the one of the *E. coli* transformed by the control plasmid are selected.

[0018]

The DNA sequence of the thus obtained mutant barnase gene may be analyzed by a conventional method, if necessary, to thereby examine the mutation in detail.

[0019]

A preferred example of the mutant barnase gene of the present invention is one which has the DNA sequence represented by SEQ ID NO:3. Compared with the nucleotide sequence of SEQ ID NO:1 coding for the wild type activity, the nucleotide sequence encoding this gene has an insertion of "T" at the 15-position from A in the initiation codon ATG and a deletion of "A" at the 333-position. According to the normal translation manner, translation will be initiated from ATG at the 1-position of this DNA sequence represented by SEQ ID NO:3, and terminated at the ninth codon which has been converted to a termination codon as a result of the insertion of the "T" at the 15-position. It is unlikely that the thus formed oligopeptide consisting of 8 amino acids has the barnase activity. Nevertheless, the barstar protein is produced by the translation of from the gene of SEQ ID NO:3. This is seemingly due to a phenomenon called "frame shift re-coding" whereby the ribosome controls the translation while automatically shifting the reading frame during the course of translation into the protein. This phenomenon, which has been reported in several genes of viruses, *E. coli* and animals. It is considered that the shifting of the reading frame suppresses translation efficiency, thereby lowering the activity of the gene.

[0020]

Moreover, there is a high possibility that there is a high possibility that a gene having a DNA sequence

derived from the DNA sequence represented by SEQ ID NO:3 by substitution, deletion, insertion or addition of one or more nucleotides will be a mutant barnase gene which provides a lowered translation efficiency when it is anther-specifically expressed in the plant. Accordingly, these genes are also included in the scope of the present invention similar to the gene of SEQ ID NO:3.

[0021]

The plant which is to be male sterilized by the mutant barnase gene is not particularly limited, so long as the gene can be transferred into the plant to provide a male sterilized plant. Examples of the plants include rice, corn, tobacco, lettuce and Brassica. Among all, rice and corn are preferred.

[0022]

In order to male sterilize a plant, the mutant barnase gene is expressed specifically in the anthers of the plant to inhibit the anther functions by the RNase activity of barnase. Such an anther-specific expression can be achieved by using the method described in WO92/13957. In brief, the mutant barnase gene is ligated to the downstream of an anther-specific promoter and then incorporated into plant cells with the use of an expression vector.

[0023]

The incorporation may be performed by using the Agrobacterium method, the electroporation method, the particle gun method, etc.

[0024]

From the transformed plant cells, a complete plant can be formed by the regeneration from a callus of the transformed plant cells in accordance with, for example, the method reported in Y. Hiel et al., Plant J. 6, 271-282:1994.

[0025]

The male sterility of the thus constructed plant can be confirmed by the inability of the plant to fertilize unless it is pollinated from another fertilizable plant.

[0026]

[Examples]

Example 1: Preparation of mutant barnase gene

Low fidelity PCR

Primer 1 (CGTTCGGCTC GATGGTACCG GTTATCAACA CGTTTGA: SEQ ID NO:6) and Primer 2 (CCTCTAGATT ATCTGATTTT TGTAAGGTC TGATAATG: SEQ ID NO:7) were synthesized by a DNA synthesizer (manufactured by Applied Biosystems) in accordance with the method described in S.L. Beaucage et al., Tetrahedron Lett., 22. 1859-1862:1982. Known plasmid pVE108 (WO92/13956) was used as a template and PCR was carried out with the use of the combination of Primers 1 and 2. SEQ ID NO:1 shows the sequence of the coding region of the barnase gene which is contained in pVE108. The reaction was performed for 50 cycles with each cycle consisting of 1 minute at 94°C, 1 minute at 57°C and 1 minute at 72°C in 10 mM Tris-HCl (pH 9.5), 50 mM KCl, 2 mM MgCl₂, and 1 mM each of dNTP, 10 ng of the template DNA

and 0.5 U of Taq DNA polymerase.

[0027]

After the completion of the reaction, the amplification product was separated by electrophoreses on agarose gel (2% SeaKem GTG agarose, 1xTAE) and purified by the DEAE-cellulose method (M. Muramatsu, Labo-manual Idenshikogaku (Gene Engineering Labo-Manual), Maruzen, pp.111:1988). By using the purified product thus obtained as a template, PCR was performed again under the conditions as defined above.

[0028]

Ligation of mutated barnase gene fragment into plasmid vector

The reaction product from the above step was digested with SacI and XbaI in a conventional manner and then purified by agarose gel electrophoresis to give an "insertion fragment". A plasmid to be used for transferring this insertion fragment into *E. coli* can be appropriately selected. For example, use may be made of plasmid pHM1 therefor. The plasmid pHM1 was constructed as follows. Plasmid pBR322 was cleaved at the EcoRI site and blunt-ended by using T4 DNA polymerase (manufactured by Takara Shuzo). Separately, lacZ expression cassette (322 bp) was excised from pUC18 with PvuII and the ends thereof were blunted. The cassette was integrated into the restriction site of the blunt-ended plasmid pBR 322 to give plasmid pHM1.

[0029]

Plasmid pHM1 was digested with SacI and XbaI followed by dephosphorylation with calf intestine alkaline phosphatase (manufactured by Takara Shuzo) to give a restriction enzyme-treated plasmid fragment. Into this restriction enzyme-treated plasmid pHM1, the above "insertion fragment" containing a barnase gene was ligated by using Takara Ligation Kit ver.1 (manufactured by Takara Shuzo).

[0030]

Introduction into *E. coli* and selection of barnase-active clone

The *E. coli* having the barnase gene introduced thereinto may be selected by, for example, the following method. Since mRNA is degraded in cells by the barnase activity, the synthesis of protein is suppressed and, in its turn, the growth of *E. coli* is inhibited. Thus, the *E. coli* transformed by the barnase gene can be selected by taking advantage of the fact that the desired transformant colony has a smaller size than the *E. coli* colony transformed by the control plasmid free from the mutant barnase gene. When the wild type barnase or a mutant barnase still maintaining the activity comparable thereto is integrated into pHM, *E. coli* cannot form any colony. Thus, clones having a sufficiently weakened barnase activity can be exclusively selected.

[0031]

Thus, the plasmid having the mutant barnase gene ligated thereto was precipitated from ethanol and then

introduced into *E. coli* LE392 strain by the electroporation method with the use of GenePulser (BioRad) in accordance with the manufacture's instruction. Also, the control plasmid free from mutant barnase was introduced into the *E. coli* LE392 strain in the same manner.

[0032]

Next, these *E. coli* transformants were plated onto an LB agar medium containing tetracycline (25 µg/ml) and incubated at room temperature (25°C) for 72 hours. Then colonies of a smaller size than the colony of *E. coli* having the control plasmid pHM1 introduced thereinto were screened as colonies of the *E. coli* transformants having the mutant barnase transferred thereinto (Table 1, A).

[0033]

Selection of mutant barnase clone by barstar gene

As described above, barstar is an enzyme acting as an antagonist to barnase. In *E. coli* wherein barstar is expressed, the enzymatic activity of barnase is inhibited and thus the degradation of mRNA, which otherwise takes place in the presence of barnase, can be inhibited. Therefore, if *E. coli* expressing barstar is transformed by a plasmid containing a barnase gene, the growth thereof will not be inhibited. Accordingly, it is expected that the thus obtained *E. coli* transformant will show little difference in growth rate from the transformant constructed by using the control plasmid free from a barnase gene and, therefore, these transformants will show

little difference in colony size too.

[0034]

An *E. coli* transformant which expressed barstar was constructed in the following manner. A barstar gene described in R.W. Hartley, J. Mol. Biol. 202, 913-915:1998 was excised with HindIII and XbaI and ligated in frame to a tac promoter (de Boer et al., Proc. Natl. Acad. Sci. USA 80, 21-25:1983). The gene, together with a chloramphenicol tolerance gene (N.K. Alton and D. Vapnek, Nature 282, 864-869:1979), was ligated into the defective transposon which lacked the transferase in a vector described in Herrero et al., J. Bacteriol. 172, 6557-6567:1990. Subsequently, the thus obtained plasmid was introduced into *E. coli* MC1061 strain thereby to transfer the transposon, which contained the barstar gene cassette, into the chromosome of the *E. coli*. It was confirmed based on the chloramphenicol tolerance that this *E. coli* was able to maintain the barstar gene cassette in a stable state. Since the *E. coli* MC1061 strain lacked the lacI gene, the tac promoter was continuously active and thus the barstar gene was constitutively expressed.

[0035]

The plasmid containing the mutant barnase gene as the insertion fragment and the control plasmid free from the same were each introduced into the above *E. coli* transformant by the electroporation method. Then the *E. coli* transformants were plated onto an LB agar medium containing IPTG (1mM) and tetracycline (25 µg/ml) and

incubated at 25°C for 72 hours. Among the colonies of the *E. coli* transformed by the mutant barnase gene, a colony being comparable in size to the colony of the *E. coli* having the control plasmid pHM1, which had been simultaneously plate-cultured, was screened and named "#4-31" (Table 1, B).

[0036]

[Table 1]

Screened clone and colony size (mm)

	Screened clone	pHM1 (control)
A: <i>E. coli</i> LE392 ^{*1}	1.77 ± 0.33	2.65 ± 0.47
B: <i>E. coli</i> with barstar gene expression ^{*2}	1.39 ± 0.10	1.55 ± 0.08

*1: Colony size after incubation on the LB agar plate containing 25 µg/mL of tetracycline at 28°C for 48 hours.

*2: Colony size after incubation on the LB agar plate containing 25 µg/mL of tetracycline at 28°C for 24 hours.

Example 2: Determination of nucleotide sequence of mutant barnase gene

From the *E. coli* screened in Example 1, a cloned mutant barnase gene appropriate for the object of the present invention was prepared. First, the clone #4-31 screened in Example 1 was propagated. Next, the mutant barnase gene fragment in the clone #4-31 was excised with KpnI and XbaI and ligated into the KpnI, XbaI-site of

pUC119 having been cleaved similarly with KpnI and XbaI. After amplifying the obtained plasmid by using *E. coli*, a reaction was performed by the cycle sequence method with the use of Taq polymerase (Taq Dye Terminator Cycle Sequencing Kit, manufactured by Applied Biosystems Inc.) in accordance with the manufacturer's protocol. Subsequently, the sequence was analyzed with a DNA Sequencer (Model 373A, manufactured by Applied Biosystems Inc.), thereby yielding the sequence represented by SEQ ID NO:3. Compared with the DNA sequence of the wild type barnase gene, this sequence had an insertion of T at the 15-position from A of the initiation codon ATG and a deletion of A at the 333-position.

[0037]

Example 3: Construction of male sterile rice by using attenuated mutant barnase gene

The attenuated mutant barnase gene integrated into pUC119 as described above was cleaved with restriction enzymes XbaI and KpnI and thus the plasmid vector pTS431 represented by SEQ ID NO:5 was constructed. Although this plasmid pTS431 differs from the known plasmid pVE108 (PCT International Publication WO92/13956) in the following points, it is substantially equivalent thereto except for the moieties of the anther-specific promoter and the mutant barnase.

[0038]

(1) In the plasmid vector pTS431 according to the present invention, the barnase gene (SEQ ID NO:1) in the

known plasmid pVE108 has been converted into the mutant barnase gene (SEQ ID NO:3).

[0039]

(2) A tobacco-origin anther specific promoter is used in the known plasmid pVE108, while the anther specific promoter of rice E1 gene (PCT International Publication WO92/13956) is used in the plasmid vector pTS431 according to the present invention.

[0040]

(3) In the present invention, a 35S3 promoter (EP 0344029) of 1376 bp, which is not used in the plasmid pVE108, is employed upstream of the barnase gene.

[0041]

(4) In the present invention, a sequence originating in the downstream of Agrobacterium T-DNA gene 7 excised from pJD884 (PCT International Publication WO93/09218) is employed whereas such a sequence is not used in the plasmid pVE108.

[0042]

(5) In the present invention, the region corresponding to lacZ has been deleted from the pUC19-origin moiety. In addition a plasmid (pTS172) having the wild type barnase gene integrated therein is represented by SEQ ID NO:4.

[0043]

The fragment of about 4.5 kbp was excised with a restriction enzyme EcoRI respectively from pTS431 (a plasmid having the mutant barnase gene transferred thereinto; SEQ ID NO:5) and pTS172 (a plasmid having the

barnase gene transferred thereinto; SEQ ID NO:4) and inserted into the EcoRI-site of the intermediate vector pSB11 (T. Komari et al., Plant J. 10(1), 165-174:1996), and further, the T-DNA region thereof was integrated into the acceptor vector pSB1 (T. Komari et al., Plant J. 10(1), 165-174:1996) by way of homologous recombination. By using *Agrobacterium tumefaciens* LBA4404 having the obtained recombinant plasmids (pSB1431 and pSB1172 respectively), rice (variety: Asanohikari) was transformed. While the transformation was performed basically in accordance with the method of Hiei et al. (Plant J. 6(2), 271-282:1994), phosphinothricine (concentration: 10 mg/L) was employed for screening transformants since the thus constructed male sterility genes contained bar gene encoding phosphinothricine acetyl transferase as a selective marker. Phosphinothricine facilitates selection of calluses having the gene transferred thereinto.

[0044]

A comparison of the rice transformant obtained with the use of the wild type barnase gene with the one obtained with the use of the mutant barnase gene indicates that the transfer of the mutant barnase gene gave rise to a remarkable improvement in the transformation efficiency and an increase in the rate of morphologically normal male sterility transformants, as shown in Table 2.

[0045]

[Table 2]

Transformation efficiency

	No. of infected calluses	No. of regenerated calluses	No. of PCR-positive lines ^{*1}	No. of morphologically normal male sterile lines
pSB1172 (control)	2838	83	52/83	9/52(17.3%)
pSB1431 (control)	787	69	43/45 ^{*2}	27/28 ^{*3} (96.4%)

*1: detection of barnase gene fraction by PCR.

*2: examined only 45 lines among 69.

*3: examined only 28 lines among 43.

[0046]

[Effects of the Invention]

In the present invention, a barnase gene having a weakened effect is constructed via mutation. A male sterility gene comprising the mutant barnase gene can be introduced into a plant to successfully produce a male sterile plant free from any unfavorable characteristic at a high efficiency with the use of a single gene without resort to the use of a barstar gene.

[0047]

[Sequence Listing]

SEQUENCE LISTING

<110> Japan Tobacco Inc.

<120> MUTANT BARNASE GENE AND TRANSGENIC PLANT TRANSFORMED BY SAID GENE

<130> 980687

<160> 7

<210> 1

<211> 343

<212> DNA

<213> Bacillus amyloliquifaciens

<220>

<221> NAME/KEY: CDS

<222> 1..336

<400> 1

atg gta ccg gtt atc aac acg ttt gac ggg gtt gcg gat tat ctt cag	48
Met Val Pro Val Ile Asn Thr Phe Asp Gly Val Ala Asp Tyr Leu Gln	
1 5 10 15	
aca tat cat aag cta cct gat aat tac att aca aaa tca gaa gca caa	96
Thr Tyr His Lys Leu Pro Asp Asn Tyr Ile Thr Lys Ser Glu Ala Gln	
20 25 30	
gcc ctc ggc tgg gtg gca tca aaa ggg aac ctt gca gac gtc gct ccg	144
Ala Leu Gly Trp Val Ala Ser Lys Gly Asn Leu Ala Asp Val Ala Pro	
35 40 45	

ggg aaa agc atc ggc gga gac atc ttc tca aac agg gaa ggc aaa ctc	192
Gly Lys Ser Ile Gly Gly Asp Ile Phe Ser Asn Arg Glu Gly Lys Leu	
50 55 60	
cgg ggc aaa agc gga cga aca tgg cgt gaa gcg gat att aac tat aca	240
Pro Gly Lys Ser Gly Arg Thr Trp Arg Glu Ala Asp Ile Asn Tyr Thr	
65 70 75 80	
tca ggc ttc aga aat tca gac cgg att ctt tac tca agc gac tgg ctg	288
Ser Gly Phe Arg Asn Ser Asp Arg Ile Leu Tyr Ser Ser Asp Trp Leu	
85 90 95	
att tac aaa aca acg gac cat tat cag acc ttt aca aaa atc aga taa	336
Ile Tyr Lys Thr Thr Asp His Tyr Gln Thr Phe Thr Lys Ile Arg	
100 105 110	
ggtaacc	343

<210> 2

<211> 112

<212> PRT

<213> Bacillus amyloliquifaciens

<400> 2

Met Val Pro Val Ile Asn Thr Phe Asp Gly Val Ala Asp Tyr Leu Gln	
1 5 10 15	
Thr Tyr His Lys Leu Pro Asp Asn Tyr Ile Thr Lys Ser Glu Ala Gln	
20 25 30	
Ala Leu Gly Trp Val Ala Ser Lys Gly Asn Leu Ala Asp Val Ala Pro	
35 40 45	
Gly Lys Ser Ile Gly Gly Asp Ile Phe Ser Asn Arg Glu Gly Lys Leu	
50 55 60	
Pro Gly Lys Ser Gly Arg Thr Trp Arg Glu Ala Asp Ile Asn Tyr Thr	

65	70	75	80
Ser Gly Phe Arg Asn Ser Asp Arg Ile Leu Tyr Ser Ser Asp Trp Leu			
	85	90	95
Ile Tyr Lys Thr Thr Asp His Tyr Gln Thr Phe Thr Lys Ile Arg			
100	105	110	

<210> 3

<211> 342

<212> DNA

<213>

<400> 3

```

atggtaaccgg ttattcaaca cgtttgacgg ggttgcggat tatcttcaga catatcataa 60
gctacctgat aattacatta caaaatcaga agcacaagcc ctcggtctggg tggcatcaaa 120
agggaaacctt gcagacgtcg ctccggggaa aagcatcggc ggagacatct tctcaaacag 180
ggaaggcaaa ctcccgggca aaagcggacg aacatggcgt gaagcggata ttaactatac 240
atcaggcttc agaaattcag accggattct ttactcaagc gactggctga ttacaaaac 300
aacggaccat taccagacct ttacaaaaat cagtaatcta ga 342

```

<210> 4

<211> 6548

<212> DNA

<213> Escherichia coli LE392

<220>

<223> Clone: pTS172

<400> 4

```

aattcaagct tgacgtcagg tggcactttt cggggaaatg tgcgcggaac ccctatttgt 60
ttatTTTTT aaatacatc aaatatgat ccgtcatga gacaataacc ctgataaatg 120

```

ctccaataat attgaaaaag gaagagtaig agtaattcaac atttccgigt cgccttatt 180
 ccccttttttg cggcattttg ccttctgtt ttgtctacc cagaaacgt ggtgaaagta 240
 aaagatgctg aagatcagtt ggggtgcacga gtgggttaca tgaactgga tctcaacagc 300
 ggtaagatcc ttgagagtt tgcctccgaa gaacgttttc caatgatgag cactttttaa 360
 gttctgtctat gtggcgcggg attatcccg attgacgccg ggcaagagca actcggctgc 420
 cgcatacact attctcagaa tgacttggtt gagtactcac cagtcacaga aaagcatctt 480
 acggatggca tgacaglaag agaattatgc agtgcgtcca taaccatgag tgataacact 540
 gcggccaact tacttctgac aacgatcgga ggaccgaagg agctaaccgc ttttttgac 600
 aacatggggg atcattgaa tgccttgat cgttgggaac cggagctgaa tgaagccata 660
 ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg caacaacgtt gcgcaacta 720
 ttaactggcg aactacttac tctagcttcc cggcaacaat taatagactg gatggaggcg 780
 gataaagttg caggaccact tctgcgtctg gcccttccgg ctggctgggt tattgctgat 840
 aaatctggag cgggtgagcg tgggtctcgc ggtatcattg cagcactggg gccagatggt 900
 aagccctccc gtatcgtagt tatctacacg acggggagtc aggcaactat ggatgaacga 960
 aatagacaga tgcctgagat aggtgcctca ctgattaage attggtaact gtcagaccaa 1020
 gtttactcat atatacttta gattgattta aaacttcatt tttaatltaa aaggatctag 1080
 gtgaagatcc tttttggctc gagtctcatg accaaaatcc cttaacgtga gttttcgttc 1140
 cactgagcgt cagaccccg agaaaagatc aaaggatctt cttagatcc ttttttctg 1200
 cgcgtaatct gctgcttgca aacaaaaaaa ccaccgtac cagcgggtgg ttgtttgccg 1260
 gatcaagagc taccactct ttttcgaag gtaactggtc tcagcagagc gcagatacca 1320
 aatactgtcc ttctagtga gccgtagttt ggccaccact tcaagaactc tgtagcaccg 1380
 cctacatacc tgcctctgct aatcctgtta ccagtggctg ctgccagtgg cgataagtcg 1440
 tgtcttaccg gggtggactc aagacgatag ttaccggata aggcgcagcg gtcgggctga 1500
 acgggggggt cgtgcacaca gccagcttg gagggaacga cctacaccga actgagatac 1560
 ctacagcgtg agcattgaga aagcgccacg ctcccgaag ggagaaaggc ggacagggtat 1620
 ccggtaagcg gcagggctcg aacaggagag cgcacgaggg agcttccagg gggaaacgcc 1680
 tggatatctt atagtccgt cgggtttcgc cactctgac ttgagcgtc atttttgtga 1740
 tgcctgtcag gggggcggag cctatggaaa aacgccagca acgcggcctt ttacgggttc 1800
 ctggcctttt gctggccttt tgcctacatg tcttttctg cgttatcccc tgattctgtg 1860

gataaccgta ttaccgectt tgagttaget galaccgctc gccgcagccg aacgaccgag 1920
cgcagcgagt cagtgagega ggaageggaa gagegccccaa tacgcaaacc gccctcccc 1980
gcgcgttggc ctgatacaga ttcatatgca cgtgttcccg atctagtaac atagatgaca 2040
cegcgcgaga taatttatcc tagtttgcgc gctatatatt gttttctalc gegtattaaa 2100
tgtataattg cgggactcta atcataaaaa cccatctcat aaataacgic atgcattaca 2160
tgttaattat tacatgctta acgtaattca acagaaatta tatgataatc atcgcaagac 2220
cggeaacagg attcaatctt aagaaacttt atlgccaaat gtttgaacga tctgcttcgg 2280
aggttacett atctgatatt tgtaaaggic tgataatggc cgttgtttt gtaaatcagc 2340
cagtcgcttg agtaaagaat ccggtctgaa ttctlgaagc ctgatgtata gttaaatcc 2400
gcttcacgcc atgttcgicc gcttttgcgc gggagtittg cttccctgtt tgagaagatg 2460
tctccgcga tcttttccc cggagcgacg tctgcaaggc tcccttttga tgcacccag 2520
ccgagggctt gtgcttctga ttttgtaatg taattatcag gtagcttatg atatgtctga 2580
agataatccg caaccccgic aaacgtgttg ataaccggta ccatcgcgac ggcttgatgg 2640
atctcttgct ggacaccggg atgctaggat gggttatcgt ggccggcgct cgtgtgtggc 2700
ttttgtaggc gccggcgacg gcggggggcaa tgtggcaggc gagtcacggc gcaagcgtgc 2760
gcaagtgact gcaacaacca aggacggtea tggcgaaagc acctcacgcg tccaccgtct 2820
acaggatgta gcagtagcac ggtgaaagaa gtgttgtccc gtccattagg tgcattctca 2880
ccgttggcca gaacaggacc gttcaacagt taggttgagt gtaggacttt tacgtggtta 2940
atgtatggca aatagtagta aattttgccc ccattggctt ggctgagata gaacatatc 3000
tggaaaagcct ctagcataat ttttttgaca gctaaacttt gcttcttgcc ttcttggtct 3060
agcaatgacg ttgcccattg cgtggcaaac atctggtaag gtaactgtat tegtgtgttc 3120
cttcaacgg ctcaatcccc acaggccaag ctatccttcc cttggcagta taggtcctt 3180
gagagattat actaccattt ttaagtgtt ataaagacga tgctctctaa ccagatcgat 3240
cagaaacaca aagtlltagc agcgtaatat cccacacaca tacacacacg aagctatgcc 3300
tctcatattt ccgagagatt ctgacagtga ccagaatgtc agaatgccat ttcattggca 3360
caagtcgac cacaagcttc ttggiggagg tcaagggtgt ctattattat tegtcttcta 3420
ggaaattatt cagaattagt gccctttatc ataactctc tctgagccga tgtgggtttg 3480
gatttcattg ttgggagcta tgcagttgcg gatattctgc tgtggaagaa caggaactta 3540
tctgcggggg tcttctgtgg ggcaacattg atatggctcc tgttcgatgt agtagaatac 3600

aatataattc cgtctcttgg ccagattgcc attcttgcca tgccttgat cticatttgg 3660
tcaaatgccg caccactctt ggacagggtat tagctttatt tcctgtggag atggtagaaa 3720
actcagctta cagaaatggc atttcacgta gtataacgca agacattagg tactaaaact 3780
caactaactg tttecgaaatt tcagggeccc tccaaggatc ccagaaatca tcctctctga 3840
acatgccttc agagaaatgg catlgaccgt ccattacaaa ctaacgtaca ctgtatctgt 3900
tctttacgac attgcatgtg gaaaggatct gaagagattt ctccctgtac ataataatct 3960
actcctttgc tacgttaata agagatgtaa aaacatgcaa cagttccagt gccaacattg 4020
tccaaggatt gtgcaattct ttctggagcg ctaaaattga ccagattaga cgcatcagaa 4080
tattgaattg cagagttagc caataatcct cataatgtta atgtgctatt gttgttcaact 4140
actcaatata gtctctggact aacaatcaga ttgtttatga tattaagggtg gttggatctc 4200
tatttggtatt gtctggcgatt ggaagttctt gcagcttgac aagtctacta tataattggt 4260
ggtattccag ataaatatata aattttaata aaacaatcac acagaaggat ctgcggccgc 4320
tagcctagge ccggggccac aaaaatctga gcttaacage acagttgctc ctctcagagc 4380
agaatcgggt attcaacacc ctcatatcaa ctactacgtt gtgtataacg gtccacatgc 4440
cggtatatac gatgactggg gtgtgtacaaa ggccggcaaca aacggcgctc ccggagttgc 4500
acacaagaaa ttgtccacta ttacagagge aagagcagca gctgacgcgt acacaacaag 4560
tcagcaaaaca gacagggtga acttcatccc caaaggagaa gctcaactca agcccaagag 4620
ctttgctaag gccctaacaa gccaccacaaa gcaaaaagcc caetggctca cgctaggaac 4680
caaaaggccc agcagtgatc cagcccaaaa agagatctcc ttgtccccgg agattacaat 4740
ggacgatttc ctctatcttt acgatctagg aaggaagttc gaaggtagaag gtgacgacac 4800
tatgttcacc actgataatg agaagggttag cctcttcaat ttacagaaaga atgttgaccc 4860
acagatgggt agagaggcct acgcagcagg tctcatcaag acgatctacc cgagtaacaa 4920
tctccaggag atcaaatacc ttcccaagaa ggttaaagat gcagtcacaa gattcaggac 4980
taattgcate aagaacacag agaaagacat atttctcaag atcagaagta ctattccagt 5040
atggacgatt caaggcttgc ttcatataacc aaggcaagta atagagattg gagtctctaa 5100
aaaggtagtt cctactgaat ctaaggccat gcatggagtc taagattcaa atcgaggatc 5160
taacagaact cgcctgaag actggcgaa acgttcataca gagcttctta cgactcaatg 5220
acaagaagaa aatcttcgtc aacatgggtg agcacgacac tcgtggtctac tccaaaaatg 5280
tcaaagatac agtctcagaa gaccaaaggg ctattgagac ttctcaacaa aggataattt 5340

cgaggaaacct cctcggattc catlgcccag ctatcigtca cttcatcgaa aggacagtag 5400
 aaaaggaagg tggctectac aaatgccatc atigcgataa aggaaaggct atcattcaag 5460
 atgectctgc cgacagtggc cccaaagatg gacccccacc cagcaggagc atcgtggaaa 5520
 aagaagacgt tccaaccacg tcttcaaagc aagtggattg atgtgacatc tccactgacg 5580
 taagggatga cgcacaatcc cactatcctt cgcaagaccc ttcctctata taaggaagtt 5640
 catttcaattt ggagaggaca cgtcgaaatc accagtctct ctctataaat ctatctctct 5700
 ctctataacc atggaccacg aacgacgccc ggcgcacatc cgccgtgcca ccgaggcgga 5760
 catgcgggag gctctgacca tcgtcaacca ctacatcgag acaagcacgg tcaacttccg 5820
 taccgagccg caggaaccgc aggagtggac ggacgacctc gtccgtctgc gggagcgcta 5880
 tccctggctc gtgcgcgagg tggacggcga ggctgcgggc atcgccctac cgggccccctg 5940
 gaaggcacgc aacgcctacg actggacggc cgagtcgacc gtgtacgtct ccccccgcca 6000
 ccagcggacg ggactgggct ccacgctcta caccacctg ctgaagtccc tggaggcaca 6060
 gggcttcaag agcgtggctc ctgcatcggt gctgccccac gacccgagcg tgcgcatgca 6120
 cgaggcgctc ggatatgccc ccgcggcat gctgcgggag gccggcttca agcacgggaa 6180
 ctggcatgac gtgggtttct ggcagctgga cttcagcctg ccggtaccgc ccgctccggt 6240
 cctgcccgtc accgagatct gagatcacgc gttctaggat ccccgatga gctaagctag 6300
 ctatatcacc aatttatgta ttacacataa tategcactc agtctttcat ctacggcaat 6360
 gtaccagctg atataatcag ttattgaaat atttctgaat ttaaacttgc atcaataaat 6420
 ttatgttttt gcttggacta taatacctga ctgtttatct tatcaataaa tatttaaaact 6480
 atatttcttt caagatggga attaacatct acaaattgcc ttttcttate gaccatgtac 6540
 gtatcgcg 6548

<210> 5

<211> 6539

<212> DNA

<213> Escherichia coli LE392

<220>

<223> Clone: pTS431

<400> 5

aattcaagct tgacgtcagg tggcactttt cggggaaaatg tgcgcggaac cectatttgt 60
ttatitttict aaatacattc aaatatgtat ccgctcatga gacaataacc ctgataaatg 120
cttcaataat attgaaaaag gaagagtatg agtattcaac atttccgtgt cgcccttatt 180
cccttttttg cggcattttg ccttctctgtt tttgctcacc cagaaacgct ggtgaaagta 240
aaagaigctg aagatcagtt ggggtgcaega gtgggttaca tcgaactgga tctcaacagc 300
ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc caatgatgag cactttttaa 360
gttctgtctat gtggcgcggtt attatcccgt attgacgccg ggcaagagca actcggtcgc 420
cgcatacact attctcagaa tgacttgggt gagtactcac cagtcacaga aaagcatctt 480
acggatggca tgacagtaag agaattatgc agtgcgtcca taaccatgag tgataacact 540
gcggccaact tacttctgac aacgatcgga ggaccgaagg agctaaccgc ttttttgcac 600
aacatggggg atcatgtaac tcgecttgat cgttgggaac cggagctgaa tgaagccata 660
ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg caacaacgtt gcgcaaaacta 720
ttaactggcg aactacttac tctagcttcc cggcaacaat taatagactg gatggaggcg 780
gataaagltg caggaccact tctgcgtctg gcccttccgg ctggctgggt tattgtgat 840
aaatctggag cgggtgagcg tgggtctcgc ggtatcattg cagcactggg gccagatggt 900
aagccctccc gtatcgtagt tatctacacg acggggagtc aggcaactat ggatgaacga 960
aatagacaga tcgctgagat aggtgcctca ctgattaage attggtaact gtcagacca 1020
gtttactcat atatacttta gattgattta aaacttcatt ttttaattta aaggatctag 1080
gtgaagatcc tttttggctc gagtctcatg accaaaatcc cttaacgtga gttttcgttc 1140
cactgagcgt cagaccccggt agaaaagatc aaaggatctt cttagatcc tttttttctg 1200
cgcgtaatct gctgcttgca aacaaaaaaa ccaccgtac cagcgggtgt ttgtttgccg 1260
gatcaagagc taccaactct ttttccgaag gtaactggct tcagcagagc gcagatacca 1320
aatactgtcc ttctagtgtg gccgtagtta ggccaccact tcaagaactc tgtagcaccg 1380
ctacataacc tcgctctgct aatctgttta ccagtggtct ctgccagttg cgataagtcg 1440
tgtcttaccg ggttggactc aagacgatag ttaccggata aggcgcagcg gtcgggctga 1500
acgggggggt cgtgcacaca gccagcttg gagcgaacga cctacaccga actgagatac 1560
ctacagcgtg agcatlgaga aagcgccacg ctccccgaag ggagaaaggc ggacaggtat 1620
ccggtlaagc gcagggtcgg aacaggagag cgcacgaggg agcttccagg gggaaacgcc 1680

tggatatttt atagtcctgt cgggttttgc cacccttgac ttgagcgctg atttttgtga 1740
 tgcctgtcag gggggcggag cctatggaaa aacgccagca acgcggcctt tttaacggtt 1800
 ctggcccttt gctggccctt tgcctacatg ttctttcctg cgtttatccc tgattctgtg 1860
 gataaccgta ttaccgcctt tgagttagct gataccgtc gccgcagccg aaagaccgag 1920
 cgcagcgagt cagttagcga ggaagcggaa gagegcceaa taagcaaacc gcctctcccc 1980
 gcgcgttggc ctgactcaga ttcttcccca tctagtaaca tagatgacac cgcgcgcgat 2040
 aattttacct agtttgcgcg ctataatttg tttctatcg cgtattaaat gtataattgc 2100
 gggactctaa tcataaaaaa ccatctcata aataacgtca tgcattacat gttaattatt 2160
 acatgcttaa cgtaattcaa cagaaattat atgataatca tcgcaagacc ggcaacagga 2220
 ttcaatctta agaaacttta ttgccaaatg ttigaacgat ctgcttcgga tctcttagat 2280
 tactgatttt tgtaaaggte tgataatggt ccgttgtttt gtaaatcagc cagtcgcttg 2340
 agtaaagaat ccggtctgaa ttcttgaagc ctgatgtata gttaatatcc gcttcacgcc 2400
 atgttctgct gcttttgcgc gggagtttgc ctccctgtt tgagaagatg tctccgccga 2460
 tgcctttccc cggagcgacg tctgcaaggt tcccttttga tgccaccag ccgagggcct 2520
 gtgcttctga ttttgtaatg taattatcag gtagcttatg atatgtctga agataatccg 2580
 caaccccgtc aaacgtgttg aataaccggt accatcgaga cggcttgatg gatctcttgc 2640
 tggacaccgg gatgctagga tgggttatcg tggccggcgt gcgtgtgtgg cttttgtagg 2700
 cgcggcgac ggcgggggca atgtggcagg tgagtcacgg tgcaagcgtg cgcaagtga 2760
 tgcaacaacc aaggacggte atggcgaaag caccctacgc gtccaccgtc tacaggatgt 2820
 agcagtagca cggtgaaaga agtgtgttgc cgtccattag gtgcattctc accgttggcc 2880
 agaacaggac cgttcaacag ttaggttgag tgiaggactt ttacgtgggt aatgtatggc 2940
 aaatagtagt aaattttgcc cccattggte tggettagat agaacatatt ctggaaagcc 3000
 tctagcatat cttttttgac agctaaactt tgcctcttgc ctcttggte tagcaatgac 3060
 gtigcccatg tctgggcaaa catctggtaa ggtaactgta ttctgttgtt ccttcaacg 3120
 gctcaatccc cacaggccaa gctatccctt ccttggcagt ataggctcct tgagagatta 3180
 tactaccatt tttaagtgt taaaaagacg atgtctctta accagatcga tcagaaacac 3240
 aaagttttag cagcgtata tccacacac atacacacac gaagctatgc cctctcattt 3300
 tccgagagat tctgacagtg accagaatgt cagaatgcca tticatgggc acaagtcgat 3360
 ccacaagctt ctgtgtggag gtaagggtg gctattattt ttctcttct aggaaattat 3420

tcagaattag tgccttttat cataacttct ctctgagccg atgtggtttt ggatttcatt 3480
 gtitgggagct atgcagttgc ggataattctg ctgtggaaga acaggaacctt atctgcgggg 3540
 gtecttgcctg gggcaacatt gatattggtc ctgttcgatg tagtagaata caatataatt 3600
 ccgtctcttt gccagattgc cattcttgcg atgcttgtga tcttcatttg gtcaaagcc 3660
 gcaccactct tggacaggta ttagctttat ttctgttgga gatggtagaa aactcagctt 3720
 acagaaatgg catttcacgt agtataacgc aagacattag gtactaaaac tcaactaact 3780
 gtittccgaat ttcaggggcc ctccaaggat cccagaaatc atcatctctg aacatgcctt 3840
 cagagaaatg gcattgaccg tccattacaa actaacgtac actgtatctg ttctttacga 3900
 cattgcatgt ggaaaggatc tgaagagatt tctcttggtt cataataatc tactcctttg 3960
 ctacgttaat aagagatgta aaaacatgca acagttccag tgccaacatt gtccaaggat 4020
 tgtgcaattc tttctggagc gctaaaattg accagattag acgcattcaga alattgaatt 4080
 gcagagttag ccaataatcc tcataatgtt aatgtgctat tgttgttcac tactcaatat 4140
 agttctggac taacaatcag attgtttatg atattaaggt gggttggtct ctattggtat 4200
 tgtcggcgat tggaaagtct tgcagcttga caagtctact atatatgtgt aggtattcca 4260
 gataaatatt aaattttaat aaaacaatca cacagaagga tctgcggccg ctagcctagg 4320
 cccgggceca caaaaatctg agcttaacag cacagttgct cctctcagag cagaatcggg 4380
 tattcaacac cctcatatca actactacgt tgtgtataac ggccacatg ccggtatata 4440
 cgatgactgg ggttgtlaca aggcggcaac aaacggcgtt cccggagtgt cacacaagaa 4500
 atttgcact attacagagg caagagcagc agctgacgcg tacacaacaa gtcagcaaac 4560
 agacaggttg aacttcattc ccaaaggaga agctcaactc aagcccaaga gctttgctaa 4620
 ggccctaaca agcccacca agcaaaaagc ccactggctc acgctaggaa ccaaaaggcc 4680
 cagcagtgat ccagcccaa aagagatctc ctttgcctcg gagattacaa tggacgattt 4740
 cctctatctt tacgactag gaaggaagtt cgaagggtgaa ggtagcgaca ctatgttcac 4800
 cactgataat gagaaggta gcctcttcaa tticagaaag aatgctgacc cacagatggt 4860
 tagagagggc tacgcagcag gtctcatcaa gacgatctac ccgagtaaca atctccagga 4920
 gatcaaatc ctccccaga aggttaaaga tgcagtcaaa agattcagga ctaattgcat 4980
 caagaacaca gagaaagaca tatttctcaa gatcagaagt actattccag tatggacgat 5040
 tcaaggttg ctccataaac caaggcaagt aatagagatt ggagtcctta aaaaggtagt 5100
 tctactgaa tctaaggcca tgcattggagt ctaagattca aatcgaggat ctaacagaac 5160

tcgccgtgaa gactggcgaa cagttcatac agagtcctttt acgactcaat gacaagaaga 5220
 aaatcttcgt caacatggig gagcaegaca ctctgggtcta ctccaaaaat gtcaaagata 5280
 cagtctcaga agaccaaaagg gctatigaga cttttcaaca aaggataatt tcgggaaacc 5340
 tcctcggatt ccattgceca gctatctgtc acttcatcga aaggacagta gaaaaggaag 5400
 gtggctccta caaatgceat catigcgata aaggaaagge taitattcaa gatgcctctg 5460
 ccgacagtgg tcccaaagat ggacccccac ccacgaggag catcgtggaa aaagaagacg 5520
 itccaaccac gtcttcaaag caagtggatt gatgtgacat ctccactgac gtaagggaatg 5580
 acgcacaatc ccactatcct tcgcaagacc ctctctctat ataaggaagt tcatttcatt 5640
 tggagaggac acgctgaaat caccagtctc tctctataaa tctatctctc tctctataac 5700
 catggacceca gaacgaagcc cgcccgacat ccgccgtgcc accgaggcgg acatgccggc 5760
 ggtctgcacc atcgtcaacc actacatcga gacaagcag gtcaacttcc gtaccgagcc 5820
 gcaggaaccg caggagtggg cggacgacct cgtccgtctg cgggagcgt atccctggct 5880
 cgtcgcgag gtggacggcg aggtcgcgg catcgctac gggggccct ggaaggcacg 5940
 caacgcctac gactggacgg ccgagtcgac cgtgtacgtc tcccccgcc accagcggac 6000
 gggactgggc tccacgtct acacccacct gctgaagtc ctggaggcac agggttcaa 6060
 gagcgtggtc gctgtcatcg ggtgcecaa cgaccgagc gtgcgcatgc acgagcgcct 6120
 cggatatgcc ccccgcgga tgetgegggc ggccggcttc aagcacggga actggcatga 6180
 cgtgggtttc tggcagctgg acttcagcct gccggtaccg ccccgctcgg tctgcccgt 6240
 caccgagatc tgagatcacg cgttctagga tccccgatg agctaagcta gctatatcat 6300
 caatttatgt attacacata atatcgcaat cagtctttca tctacggcaa tgtaccagct 6360
 gatataatca gttattgaaa tttttctgaa tttaaacttg catcaataaa tttatgtttt 6420
 tgcttggact ataatacctg acttgttatt ttatcaataa atatttaaac tatatttctt 6480
 tcaagatggg aattaacatc tacaaattgc ctttcttat cgaccatgta cgtatcgcg 6539

<210> 6

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer 1

<400> 6

cgttcggctc gatgggaccg gttatcaaca cgtttga 37

<210> 7

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer 2

<400> 7

cctctagatt atctgatttt tgtaaaggtc tgataatg 38

[Name of Document] Abstract

[Abstract]

 [Problem] The present invention aims at constructing a male sterile plant free from any undesirable characteristic by anther-specifically expressing barnase gene alone.

 [Means for Solving] A barnase gene, which has a mutation and thus sustains its activity at a lowered level, is transferred into a plant and anther-specifically expressed, thereby efficiently providing a male sterile transgenic plant.

[Selected Drawing] None